



0 - 0 - 1

## GUARANTEED ANALYSIS

Soluble Potash ( $K_2O$ ) 1%

Derived from seaweed extract (*Ascophyllum nodosum*) and potassium hydroxide

### F000148

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

For your convenience, TOGGLE® product information is available on Agrian database (<http://agrian.com/home/>) as well as in Crop Data Management Systems database (<http://www.cdms.net/>)

**STORAGE AND HANDLING:** This product should be stored away from intense sunlight and heat.

### STIR BEFORE USE

### CAUTION

Keep out of reach of children.

### WARRANTY STATEMENT:

The manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of the manufacturer. In no case shall the manufacturer be liable for consequential, special or indirect damages resulting from the use or handling of this product. The manufacturer makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

### Manufactured and guaranteed by:

Acadian Seaplants Limited  
30 Brown Avenue, Dartmouth, Nova Scotia  
Canada, B3B 1X8  
Tel: +1 902 468 2840  
Fax: +1 902 468 3474

Acadian Plant Health™ is a division of  
Acadian Seaplants Limited



**NET CONTENT:**  
2.5 U.S. Gal.  
9.4 Liters

**NET WEIGHT:**  
21.53 lb  
9.77 kg

USA\_TG1427a-2\_0623



**ACADIAN™**  
PLANT HEALTH

**GENERAL INFORMATION:** Consistent use of **TOGGLE**® supplements a well-balanced crop nutrition program. Use **TOGGLE**® to:

- Maximize crop potential during periods of stress
- Improve drought tolerance and resistance to extreme temperatures (cold and heat)
- Increase and maximize yield
- Optimize nutrient uptake
- Improve root growth and development
- Improve photosynthesis

To achieve the desired results, the levels of major and minor nutrients must be adequate to support the increase in production. The following rates and timings of application are recommended for optimum efficacy.

**COMPATIBILITY:** **TOGGLE**® is compatible with most insecticides, fungicides and fertilizers. When mixing with calcium products, thoroughly mix **TOGGLE**® with the water in the tank prior to adding the calcium product. If interaction of chemicals is unknown, a “jar” compatibility test is suggested.

## DIRECTIONS FOR USE

**TOGGLE**® fully dissolves in water and is suitable for use in liquid foliar, soil applied, and irrigation water applications. With the versatility to apply in-furrow or by foliar application, **TOGGLE**® works synergistically with a grower’s fertilizer program to improve the crop’s use of nutrients and make the best use of production inputs. Regular applications are important for maximizing crop potential during unexpected stress.

**TOGGLE**® can be tank mixed with most commonly used insecticides, fungicides and herbicides.

The product may not be applied near water, storm drains or drainage ditches or to any impervious surface.

The product may not be applied if heavy rain is expected.

The product may only be applied to the intended application site.

Crop	Application	Rate per Acre	Recommendation and Timings
Corn (Field corn, silage, sweet corn, popcorn, seed corn)	1st	8 ounces	In-furrow or banded at planting, or with row starter. †
	2nd	32 to 48 ounces	Foliar broadcast spray at V4 - VT. Note: Always follow fungicide labels regarding surfactant use at/near tassel stages.
Soybean	1st	8 ounces	In-furrow or banded at planting, or with row starter. †
	2nd	32 to 48 ounces	Foliar broadcast from V4 - R3.
Wheat	1st	8 ounces	In-furrow or banded at planting, or with row starter. †
	2nd	32 to 48 ounces	At tiller initiation.
	3rd	32 to 48 ounces	At early boot to early panicle emergence.
Cotton	1st	8 ounces	In-furrow or banded at planting, or with row starter. †
	2nd	32 to 48 ounces	Foliar broadcast at pinhead square. †
	3rd	32 to 48 ounces	Foliar broadcast two weeks after pinhead square.
Rice	1st	8 ounces	In-furrow or banded at planting, or with row starter. †
	2nd	32 to 48 ounces	Foliar broadcast at tiller initiation. †
	3rd	32 to 48 ounces	Foliar broadcast at early panicle emergence. †
Alfalfa, Mixed Hay	1st	8 ounces	In-furrow or banded at planting, or with row starter. †
	2nd	32 to 48 ounces	Soon after each cutting. †
	3rd	32 to 48 ounces	Two weeks after cutting. †
Potatoes	1st	8 ounces	In-furrow or banded at planting, or with row starter. †
	2nd	32 to 48 ounces	At tuber initiation. †
	3rd	32 to 48 ounces	Through late tuber set (if more potatoes are desired). †
Peanuts	1st	8 ounces	In-furrow or banded at planting, or with row starter. †
	2nd	32 to 48 ounces	After plants are established through to harvest. †
All other row crops	1st	8 ounces	In-furrow or banded at planting, or with row starter. †
	2nd	32 to 48 ounces	After planting through to harvest. †

Additional applications are recommended prior to major stress events or after severe hail or insect damage or herbicide damage.

† Key timings for best results. For more information see your **TOGGLE**® authorized distributor or **TOGGLE**® representative.



0 - 0 - 1

## GUARANTEED ANALYSIS

Soluble Potash (K<sub>2</sub>O) 1%

Derived from seaweed extract (*Ascophyllum nodosum*) and potassium hydroxide

Density at 68°F (20°C) 8.67 lb / U.S. Gal (1.04 kg/L)

F000148

**STORAGE AND HANDLING:** This product should be stored away from intense sunlight and heat.

### STIR BEFORE USE

### CAUTION

Keep out of reach of children.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

For your convenience, **TOGGLE**® product information is available on Agrian database (<http://agrian.com/home/>) as well as in Crop Data Management Systems database (<http://www.cdms.net/>)

### WARRANTY STATEMENT:

The manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of the manufacturer. In no case shall the manufacturer be liable for consequential, special or indirect damages resulting from the use or handling of this product. The manufacturer makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

### Manufactured and guaranteed by:

Acadian Seaplants Limited  
30 Brown Avenue, Dartmouth, Nova Scotia  
Canada, B3B 1X8  
Tel: +1 902 468 2840  
Fax: +1 902 468 3474

Acadian Plant Health™ is a division of  
Acadian Seaplants Limited



**NET CONTENT:**  
2 x 2.5 U.S. Gal.  
2 x 9.4 Liters

**NET WEIGHT:**  
43.06 lb  
19.54 kg

**GENERAL INFORMATION:** Consistent use of **TOGGLE**® supplements a well-balanced crop nutrition program. Use **TOGGLE**® to:

- Maximize crop potential during periods of stress
- Improve drought tolerance and resistance to extreme temperatures (cold and heat)
- Increase and maximize yield
- Optimize nutrient uptake
- Improve root growth and development
- Improve photosynthesis

To achieve the desired results, the levels of major and minor nutrients must be adequate to support the increase in production. The following rates and timings of application are recommended for optimum efficacy.

**COMPATIBILITY:** **TOGGLE**® is compatible with most insecticides, fungicides and fertilizers. When mixing with calcium products, thoroughly mix **TOGGLE**® with the water in the tank prior to adding the calcium product. If interaction of chemicals is unknown, a "jar" compatibility test is suggested.

## DIRECTIONS FOR USE

**TOGGLE**® fully dissolves in water and is suitable for use in liquid foliar, soil applied, and irrigation water applications. With the versatility to apply in-furrow or by foliar application, **TOGGLE**® works synergistically with a grower's fertilizer program to improve the crop's use of nutrients and make the best use of production inputs. Regular applications are important for maximizing crop potential during unexpected stress.

**TOGGLE**® can be tank mixed with most commonly used insecticides, fungicides and herbicides.

The product may not be applied near water, storm drains or drainage ditches or to any impervious surface.

The product may not be applied if heavy rain is expected.

The product may only be applied to the intended application site.

Crop	Application	Rate per Acre	Recommendation and Timings
Corn (Field corn, silage, sweet corn, popcorn, seed corn)	1st	8 ounces	In-furrow or banded at planting, or with row starter. †
	2nd	32 to 48 ounces	Foliar broadcast spray at V4 - VT. Note: Always follow fungicide labels regarding surfactant use at/near tassel stages.
Soybean	1st	8 ounces	In-furrow or banded at planting, or with row starter. †
	2nd	32 to 48 ounces	Foliar broadcast from V4 - R3.
Wheat	1st	8 ounces	In-furrow or banded at planting, or with row starter. †
	2nd	32 to 48 ounces	At tiller initiation.
	3rd	32 to 48 ounces	At early boot to early panicle emergence.
Cotton	1st	8 ounces	In-furrow or banded at planting, or with row starter. †
	2nd	32 to 48 ounces	Foliar broadcast at pinhead square. †
	3rd	32 to 48 ounces	Foliar broadcast two weeks after pinhead square.
Rice	1st	8 ounces	In-furrow or banded at planting, or with row starter. †
	2nd	32 to 48 ounces	Foliar broadcast at tiller initiation. †
	3rd	32 to 48 ounces	Foliar broadcast at early panicle emergence. †
Alfalfa, Mixed Hay	1st	8 ounces	In-furrow or banded at planting, or with row starter. †
	2nd	32 to 48 ounces	Soon after each cutting. †
	3rd	32 to 48 ounces	Two weeks after cutting. †
Potatoes	1st	8 ounces	In-furrow or banded at planting, or with row starter. †
	2nd	32 to 48 ounces	At tuber initiation. †
	3rd	32 to 48 ounces	Through late tuber set (if more potatoes are desired). †
Peanuts	1st	8 ounces	In-furrow or banded at planting, or with row starter. †
	2nd	32 to 48 ounces	After plants are established through to harvest. †
All other row crops	1st	8 ounces	In-furrow or banded at planting, or with row starter. †
	2nd	32 to 48 ounces	After planting through to harvest. †

Additional applications are recommended prior to major stress events or after severe hail or insect damage or herbicide damage.

† Key timings for best results. For more information see your **TOGGLE**® authorized distributor or **TOGGLE**® representative.